

Distance-regular graphs, pseudo primitive idempotents, and the Terwilliger algebra

Paul Terwilliger Chih-wen Weng

Abstract

Let Γ denote a distance-regular graph with diameter $D \geq 3$, intersection numbers a_i, b_i, c_i and Bose-Mesner algebra \mathbf{M} . For $\theta \in \mathbb{C} \cup \infty$ we define a 1 dimensional subspace of \mathbf{M} which we call $\mathbf{M}(\theta)$. If $\theta \in \mathbb{C}$ then $\mathbf{M}(\theta)$ consists of those Y in \mathbf{M} such that $(A - \theta I)Y \in \mathbb{C}A_D$, where A (resp. A_D) is the adjacency matrix (resp. D th distance matrix) of Γ . If $\theta = \infty$ then $\mathbf{M}(\theta) = \mathbb{C}A_D$. By a *pseudo primitive idempotent* for θ we mean a nonzero element of $\mathbf{M}(\theta)$. We use these as follows. Let X denote the vertex set of Γ and fix $x \in X$. Let \mathbf{T} denote the subalgebra of $\text{Mat}_X(\mathbb{C})$ generated by $A, E_0^*, E_1^*, \dots, E_D^*$, where E_i^* denotes the projection onto the i th subconstituent of Γ with respect to x . \mathbf{T} is called the Terwilliger algebra. Let W denote an irreducible \mathbf{T} -module. By the *endpoint* of W we mean $\min\{i | E_i^* W \neq 0\}$. W is called *thin* whenever $\dim(E_i^* W) \leq 1$ for $0 \leq i \leq D$. Let $V = \mathbb{C}^X$ denote the standard \mathbf{T} -module. Fix $0 \neq v \in E_1^* V$ with v orthogonal to the all 1's vector. We define $(\mathbf{M}; v) := \{P \in \mathbf{M} | Pv \in E_D^* V\}$. We show the following are equivalent: (i) $\dim(\mathbf{M}; v) \geq 2$; (ii) v is contained in a thin irreducible \mathbf{T} -module with endpoint 1. Suppose (i), (ii) hold. We show $(\mathbf{M}; v)$ has a basis J, E where J has all entries 1 and E is defined as follows. Let W denote the \mathbf{T} -module which satisfies (ii). Observe $E_1^* W$ is an eigenspace for $E_1^* A E_1^*$; let η denote the corresponding eigenvalue. Define $\tilde{\eta} = -1 - b_1(1 + \eta)^{-1}$ if $\eta \neq -1$ and $\tilde{\eta} = \infty$ if $\eta = -1$. Then E is a pseudo primitive idempotent for $\tilde{\eta}$.

Keywords: distance-regular graph, pseudo primitive idempotents, subconstituent algebra, Terwilliger algebra.

AMS Subject Classification: 05E30.

1 Introduction

Let Γ denote a distance-regular graph with diameter $D \geq 3$, intersection numbers a_i, b_i, c_i , Bose-Mesner algebra \mathbf{M} and path-length distance function ∂ (see Section 2 for formal definitions). In order to state our main theorems we make a few comments. Let X denote the vertex set of Γ . Let $V = \mathbb{C}^X$ denote the vector space over \mathbb{C} consisting of column vectors whose coordinates are indexed by X and whose entries are in \mathbb{C} . We endow V with the Hermitean inner product $\langle \cdot, \cdot \rangle$ satisfying $\langle u, v \rangle = u^t \bar{v}$ for all $u, v \in V$. For each $y \in X$ let \hat{y} denote the vector in V with a 1 in the y coordinate and 0 in all other coordinates. We observe $\{\hat{y} | y \in X\}$ is an orthonormal basis for V . Fix $x \in X$. For $0 \leq i \leq D$ let E_i^* denote the diagonal matrix in $\text{Mat}_X(\mathbb{C})$ which has yy entry 1 (resp. 0) whenever $\partial(x, y) = i$ (resp. $\partial(x, y) \neq i$). We observe E_i^* acts on V as the projection onto the i th subconstituent of Γ with respect to x . For $0 \leq i \leq D$ define $s_i = \sum \hat{y}$, where the sum is over all vertices $y \in X$ such that $\partial(x, y) = i$. We observe $s_i \in E_i^*V$. Let v denote a nonzero vector in E_1^*V which is orthogonal to s_1 . We define

$$(\mathbf{M}; v) := \{P \in \mathbf{M} \mid Pv \in E_D^*V\}.$$

We observe $(\mathbf{M}; v)$ is a subspace of \mathbf{M} . We consider the dimension of $(\mathbf{M}; v)$. We first observe $(\mathbf{M}; v) \neq 0$. To see this, let J denote the matrix in $\text{Mat}_X(\mathbb{C})$ which has all entries 1. It is known J is contained in \mathbf{M} [2, p. 64]. In fact $J \in (\mathbf{M}; v)$; the reason is $Jv = 0$ since v is orthogonal to s_1 . Apparently $(\mathbf{M}; v)$ is nonzero so it has dimension at least 1. We now consider when does $(\mathbf{M}; v)$ have dimension at least 2? To answer this question we recall the Terwilliger algebra. Let \mathbf{T} denote the subalgebra of $\text{Mat}_X(\mathbb{C})$ generated by $A, E_0^*, E_1^*, \dots, E_D^*$, where A denotes the adjacency matrix of Γ . The algebra \mathbf{T} is known as the *Terwilliger algebra* (or *subconstituent algebra*) of Γ with respect to x [19, 20, 21]. By a *\mathbf{T} -module* we mean a subspace $W \subseteq V$ such that $\mathbf{T}W \subseteq W$. Let W denote a \mathbf{T} -module. We say W is *irreducible* whenever $W \neq 0$ and W does not contain a \mathbf{T} -module other than 0 and W . Let W denote an irreducible \mathbf{T} -module. By the *endpoint* of W we mean the minimal integer i ($0 \leq i \leq D$) such that $E_i^*W \neq 0$. We say W is *thin* whenever E_i^*W has dimension at most 1 for $0 \leq i \leq D$. We now state our main theorem.

Theorem 1.1. *Let v denote a nonzero vector in E_1^*V which is orthogonal to s_1 . Then the following (i), (ii) are equivalent.*

(i) $(\mathbf{M}; v)$ has dimension at least 2.

(ii) v is contained in a thin irreducible \mathbf{T} -module with endpoint 1.

Suppose (i), (ii) hold above. Then $(\mathbf{M}; v)$ has dimension exactly 2.

With reference to Theorem 1.1, suppose for the moment that (i), (ii) hold. We find a basis for $(\mathbf{M}; v)$. To describe our basis we need some notation. Let $\theta_0 > \theta_1 > \cdots > \theta_D$ denote the distinct eigenvalues of A , and for $0 \leq i \leq D$ let E_i denote the primitive idempotent of \mathbf{M} associated with θ_i . We recall E_i satisfies $(A - \theta_i I)E_i = 0$. We introduce a type of element in \mathbf{M} which generalizes the E_0, E_1, \dots, E_D . We call this type of element a *pseudo primitive idempotent* for Γ . In order to define the pseudo primitive idempotents, we first define for each $\theta \in \mathbb{C} \cup \infty$ a subspace of \mathbf{M} which we call $\mathbf{M}(\theta)$. For $\theta \in \mathbb{C}$, $\mathbf{M}(\theta)$ consists of those elements Y of \mathbf{M} such that $(A - \theta I)Y \in \mathbb{C}A_D$, where A_D is the D th distance matrix of Γ . We define $\mathbf{M}(\infty) = \mathbb{C}A_D$. We show $\mathbf{M}(\theta)$ has dimension 1 for all $\theta \in \mathbb{C} \cup \infty$. Given distinct $\theta, \theta' \in \mathbb{C} \cup \infty$, we show $\mathbf{M}(\theta) \cap \mathbf{M}(\theta') = 0$. For $0 \leq i \leq D$ we show $\mathbf{M}(\theta_i) = \mathbb{C}E_i$. Let $\theta \in \mathbb{C} \cup \infty$. By a *pseudo primitive idempotent* for θ , we mean a nonzero element of $\mathbf{M}(\theta)$. Before proceeding we define an involution on $\mathbb{C} \cup \infty$. For $\eta \in \mathbb{C} \cup \infty$ we define

$$\tilde{\eta} = \begin{cases} \infty & \text{if } \eta = -1, \\ -1 & \text{if } \eta = \infty, \\ -1 - \frac{b_1}{1+\eta} & \text{if } \eta \neq -1, \eta \neq \infty. \end{cases}$$

We observe $\tilde{\tilde{\eta}} = \eta$ for $\eta \in \mathbb{C} \cup \infty$. Let W denote a thin irreducible \mathbf{T} -module with endpoint 1. Observe E_1^*W is a one dimensional eigenspace for $E_1^*AE_1^*$; let η denote the corresponding eigenvalue. We call η the *local eigenvalue* of W .

Theorem 1.2. *Let v denote a nonzero vector in E_1^*V which is orthogonal to s_1 . Suppose v satisfies the equivalent conditions (i), (ii) in Theorem 1.1. Let W denote the \mathbf{T} -module from part (ii) of that theorem and let η denote the local eigenvalue for W . Let E denote a pseudo primitive idempotent for $\tilde{\eta}$. Then J, E form a basis for $(\mathbf{M}; v)$.*

We comment on when the scalar $\tilde{\eta}$ from Theorem 1.2 is an eigenvalue of Γ . Let W denote a thin irreducible \mathbf{T} -module with endpoint 1 and local

eigenvalue η . It is known $\tilde{\theta}_1 \leq \eta \leq \tilde{\theta}_D$ [18, Theorem 1]. If $\eta = \tilde{\theta}_1$ then $\tilde{\eta} = \theta_1$. If $\eta = \tilde{\theta}_D$ then $\tilde{\eta} = \theta_D$. We show that if $\tilde{\theta}_1 < \eta < \tilde{\theta}_D$ then $\tilde{\eta}$ is not an eigenvalue of Γ .

The paper is organized as follows. In section 2 we give some preliminaries on distance-regular graphs. In section 3 and section 4 we review some basic results on the Terwilliger algebra and its modules. We prove Theorem 1.1 in section 5. In section 6 we discuss pseudo primitive idempotents. In section 7 we discuss local eigenvalues. We prove Theorem 1.2 in section 8.

2 Preliminaries

In this section we review some definitions and basic concepts. See the books by Bannai and Ito [2] or Brouwer, Cohen, and Neumaier [4] for more background information.

Let X denote a nonempty finite set. Let $\text{Mat}_X(\mathbb{C})$ denote the \mathbb{C} -algebra consisting of all matrices whose rows and columns are indexed by X and whose entries are in \mathbb{C} . Let $V = \mathbb{C}^X$ denote the vector space over \mathbb{C} consisting of column vectors whose coordinates are indexed by X and whose entries are in \mathbb{C} . We observe $\text{Mat}_X(\mathbb{C})$ acts on V by left multiplication. We endow V with the Hermitean inner product $\langle \cdot, \cdot \rangle$ which satisfies $\langle u, v \rangle = u^t \bar{v}$ for all $u, v \in V$, where t denotes transpose and $\bar{\cdot}$ denotes complex conjugation. For all $y \in X$, let \hat{y} denote the element of V with a 1 in the y coordinate and 0 in all other coordinates. We observe $\{\hat{y} \mid y \in X\}$ is an orthonormal basis for V .

Let $\Gamma = (X, R)$ denote a finite, undirected, connected graph without loops or multiple edges, with vertex set X , edge set R , path-length distance function ∂ and diameter $D := \max\{\partial(x, y) \mid x, y \in X\}$. We say Γ is *distance-regular* whenever for all integers h, i, j ($0 \leq h, i, j \leq D$) and for all $x, y \in X$ with $\partial(x, y) = h$, the number

$$p_{ij}^h = |\{z \in X \mid \partial(x, z) = i, \partial(z, y) = j\}| \quad (2.1)$$

is independent of x and y . The integers p_{ij}^h are called the *intersection numbers* for Γ . Observe $p_{ij}^h = p_{ji}^h$ ($0 \leq h, i, j \leq D$). We abbreviate $c_i := p_{1i-1}^i$ ($1 \leq i \leq D$), $a_i := p_{1i}^i$ ($0 \leq i \leq D$), $b_i := p_{1i+1}^i$ ($0 \leq i \leq D-1$), $k_i := p_{ii}^0$

$(0 \leq i \leq D)$, and for convenience we set $c_0 := 0$ and $b_D := 0$. Note that $b_{i-1}c_i \neq 0$ ($1 \leq i \leq D$).

For the rest of this paper we assume $\Gamma = (X, R)$ is distance-regular with diameter $D \geq 3$. By (2.1) and the triangle inequality,

$$p_{i1}^h = 0 \quad \text{if } |h - i| > 1 \quad (0 \leq h, i \leq D), \quad (2.2)$$

$$p_{ij}^1 = 0 \quad \text{if } |i - j| > 1 \quad (0 \leq i, j \leq D). \quad (2.3)$$

Observe Γ is regular with valency $k = k_1 = b_0$, and that $k = c_i + a_i + b_i$ for $0 \leq i \leq D$. By [4, p. 127] we have

$$k_{i-1}b_{i-1} = k_i c_i \quad (1 \leq i \leq D). \quad (2.4)$$

We recall the Bose-Mesner algebra of Γ . For $0 \leq i \leq D$ let A_i denote the matrix in $\text{Mat}_X(\mathbb{C})$ which has yz entry

$$(A_i)_{yz} = \begin{cases} 1 & \text{if } \partial(y, z) = i \\ 0 & \text{if } \partial(y, z) \neq i \end{cases} \quad (y, z \in X).$$

We call A_i the *i th distance matrix* of Γ . For notational convenience we define $A_i = 0$ for $i < 0$ and $i > D$. Observe (ai) $A_0 = I$; (aii) $\sum_{i=0}^D A_i = J$; (aiii) $\overline{A_i} = A_i$ ($0 \leq i \leq D$); (aiv) $A_i^t = A_i$ ($0 \leq i \leq D$), (av) $A_i A_j = \sum_{h=0}^D p_{ij}^h A_h$ ($0 \leq i, j \leq D$), where I denotes the identity matrix and J denotes the all ones matrix. We abbreviate $A := A_1$ and call this the *adjacency matrix* of Γ . Let \mathbf{M} denote the subalgebra of $\text{Mat}_X(\mathbb{C})$ generated by A . Using (ai)–(av) we find A_0, A_1, \dots, A_D form a basis of \mathbf{M} . We call \mathbf{M} the *Bose-Mesner algebra* of Γ . By [2, p. 59, p. 64], \mathbf{M} has a second basis E_0, E_1, \dots, E_D such that (ei) $E_0 = |X|^{-1}J$; (eii) $\sum_{i=0}^D E_i = I$; (eiii) $\overline{E_i} = E_i$ ($0 \leq i \leq D$); (eiv) $E_i^t = E_i$ ($0 \leq i \leq D$); (ev) $E_i E_j = \delta_{ij} E_i$ ($0 \leq i, j \leq D$). We call E_0, E_1, \dots, E_D the *primitive idempotents* for Γ . Since E_0, E_1, \dots, E_D form a basis for \mathbf{M} there exists complex scalars $\theta_0, \theta_1, \dots, \theta_D$ such that $A = \sum_{i=0}^D \theta_i E_i$. By this and (ev) we find $AE_i = \theta_i E_i$ for $0 \leq i \leq D$. Using (aiii) and (eiii) we find each of $\theta_0, \theta_1, \dots, \theta_D$ is a real number. Observe $\theta_0, \theta_1, \dots, \theta_D$ are mutually distinct since A generates \mathbf{M} . By [2, p.197] we have $\theta_0 = k$ and $-k \leq \theta_i \leq k$ for $0 \leq i \leq D$. Throughout this paper, we assume E_0, E_1, \dots, E_D are indexed so that $\theta_0 > \theta_1 > \dots > \theta_D$. We call θ_i the *i th eigenvalue* of Γ .

We recall some polynomials. To motivate these we make a comment. Setting $i = 1$ in (av) and using (2.2),

$$AA_j = b_{j-1}A_{j-1} + a_jA_j + c_{j+1}A_{j+1} \quad (0 \leq j \leq D-1), \quad (2.5)$$

where $b_{-1} = 0$. Let λ denote an indeterminate and let $\mathbb{C}[\lambda]$ denote the \mathbb{C} -algebra consisting of all polynomials in λ which have coefficients in \mathbb{C} . Let f_0, f_1, \dots, f_D denote the polynomials in $\mathbb{C}[\lambda]$ which satisfy $f_0 = 1$ and

$$\lambda f_j = b_{j-1}f_{j-1} + a_jf_j + c_{j+1}f_{j+1} \quad (0 \leq j \leq D-1), \quad (2.6)$$

where $f_{-1} = 0$. For $0 \leq j \leq D$ the degree of f_j is exactly j . Comparing (2.5) and (2.6) we find $A_j = f_j(A)$.

3 The Terwilliger algebra

For the remainder of this paper we fix $x \in X$. For $0 \leq i \leq D$ let $E_i^* = E_i^*(x)$ denote the diagonal matrix in $\text{Mat}_X(\mathbb{C})$ which has yy entry

$$(E_i^*)_{yy} = \begin{cases} 1 & \text{if } \partial(x, y) = i \\ 0 & \text{if } \partial(x, y) \neq i \end{cases} \quad (y \in X). \quad (3.1)$$

We call E_i^* the i th dual idempotent of Γ with respect to x . For convenience we define $E_i^* = 0$ for $i < 0$ and $i > D$. We observe (i) $\sum_{i=0}^D E_i^* = I$; (ii) $\overline{E_i^*} = E_i^*$ ($0 \leq i \leq D$), (iii) $E_i^{*t} = E_i^*$ ($0 \leq i \leq D$), (iv) $E_i^* E_j^* = \delta_{ij} E_i^*$ ($0 \leq i, j \leq D$). The E_i^* have the following interpretation. Using (3.1) we find

$$E_i^* V = \text{span}\{\hat{y} | y \in X, \partial(x, y) = i\} \quad (0 \leq i \leq D).$$

By this and since $\{\hat{y} | y \in X\}$ is an orthonormal basis for V ,

$$V = E_0^* V + E_1^* V + \dots + E_D^* V \quad (\text{orthogonal direct sum}).$$

For $0 \leq i \leq D$, E_i^* acts on V as the projection onto $E_i^* V$. We call $E_i^* V$ the i th subconstituent of Γ with respect to x . For $0 \leq i \leq D$ we define $s_i = \sum \hat{y}$, where the sum is over all vertices $y \in X$ such that $\partial(x, y) = i$. We observe $s_i \in E_i^* V$.

Let $\mathbf{T} = \mathbf{T}(x)$ denote the subalgebra of $\text{Mat}_X(\mathbb{C})$ generated by $A, E_0^*, E_1^*, \dots, E_D^*$. The algebra \mathbf{T} is semisimple but not commutative in general [19,

Lemma 3.4]. We call \mathbf{T} the *Terwilliger algebra* (or *subconstituent algebra*) of Γ with respect to x . We refer the reader to [1, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24] for more information on the Terwilliger algebra. We will use the following facts. Pick any integers h, i, j ($0 \leq h, i, j \leq D$). By [19, Lemma 3.2] we have $E_i^* A_h E_j^* = 0$ if and only if $p_{ij}^h = 0$. By this and (2.2), (2.3) we find

$$E_i^* A_h E_1^* = 0 \quad \text{if} \quad |h - i| > 1 \quad (0 \leq h, i \leq D), \quad (3.2)$$

$$E_i^* A E_j^* = 0 \quad \text{if} \quad |i - j| > 1 \quad (0 \leq i, j \leq D). \quad (3.3)$$

Lemma 3.1. *The following (i), (ii) hold for $0 \leq i \leq D$.*

$$(i) \quad E_i^* J E_1^* = E_i^* A_{i-1} E_1^* + E_i^* A_i E_1^* + E_i^* A_{i+1} E_1^*.$$

$$(ii) \quad A_i E_1^* = E_{i-1}^* A_i E_1^* + E_i^* A_i E_1^* + E_{i+1}^* A_i E_1^*.$$

Proof. (i) Recall $J = \sum_{h=0}^D A_h$ so $E_i^* J E_1^* = \sum_{h=0}^D E_i^* A_h E_1^*$. Evaluating this using (3.2) we obtain the result.

(ii) Recall $I = \sum_{h=0}^D E_h^*$ so $A_i E_1^* = \sum_{h=0}^D E_h^* A_i E_1^*$. Evaluating this using (3.2) we obtain the result. \square

Lemma 3.2. *For $0 \leq i \leq D - 1$ we have*

$$E_{i+1}^* A_i E_1^* - E_i^* A_{i+1} E_1^* = \sum_{h=0}^i A_h E_1^* - \sum_{h=0}^i E_h^* J E_1^*. \quad (3.4)$$

Proof. Evaluate each term in the right-hand side of (3.4) using Lemma 3.1 and simplify the result. \square

Corollary 3.3. *Let v denote a vector in $E_1^* V$ which is orthogonal to s_1 . Then for $0 \leq i \leq D - 1$ we have*

$$E_{i+1}^* A_i v - E_i^* A_{i+1} v = \sum_{h=0}^i A_h v. \quad (3.5)$$

Proof. Apply all terms of (3.4) to v and evaluate the result using $E_1^* v = v$ and $Jv = 0$. \square

Lemma 3.4. *The following (i), (ii) hold for $1 \leq i \leq D - 1$.*

$$(i) \quad E_{i+1}^* A E_i^* A_{i-1} E_1^* = c_i E_{i+1}^* A_i E_1^*$$

$$(ii) \ E_{i-1}^* A E_i^* A_{i+1} E_1^* = b_i E_{i-1}^* A_i E_1^*.$$

Proof. (i) For all $y, z \in X$, on either side the yz entry is equal to c_i if $\partial(x, y) = i + 1$, $\partial(x, z) = 1$, $\partial(y, z) = i$, and zero otherwise.

(ii) For all $y, z \in X$, on either side the yz entry is equal to b_i if $\partial(x, y) = i - 1$, $\partial(x, z) = 1$, $\partial(y, z) = i$, and zero otherwise. \square

Corollary 3.5. *Let v denote a vector in $E_1^* V$. Then the following (i), (ii) hold for $1 \leq i \leq D - 1$.*

(i) *Suppose $E_i^* A_{i-1} v = 0$. Then $E_{i+1}^* A_i v = 0$.*

(ii) *Suppose $E_i^* A_{i+1} v = 0$. Then $E_{i-1}^* A_i v = 0$.*

Proof. In Lemma 3.4(i),(ii) apply both sides to v and use $E_1^* v = v$. \square

4 The modules of the Terwilliger algebra

Let \mathbf{T} denote the Terwilliger algebra of Γ with respect to x . By a \mathbf{T} -module we mean a subspace $W \subseteq V$ such that $BW \subseteq W$ for all $B \in \mathbf{T}$. Let W denote a \mathbf{T} -module. Then W is said to be *irreducible* whenever W is nonzero and W contains no \mathbf{T} -modules other than 0 and W . Let W denote an irreducible \mathbf{T} -module. Then W is the orthogonal direct sum of the nonzero spaces among $E_0^* W, E_1^* W, \dots, E_D^* W$ [19, Lemma 3.4]. By the *endpoint* of W we mean $\min\{i | 0 \leq i \leq D, E_i^* W \neq 0\}$. By the *diameter* of W we mean $|\{i | 0 \leq i \leq D, E_i^* W \neq 0\}| - 1$. We say W is *thin* whenever $E_i^* W$ has dimension at most 1 for $0 \leq i \leq D$. There exists a unique irreducible \mathbf{T} -module which has endpoint 0 [10, Prop. 8.4]. This module is called V_0 . For $0 \leq i \leq D$ the vector s_i is a basis for $E_i^* V_0$ [19, Lemma 3.6]. Therefore V_0 is thin with diameter D . The module V_0 is orthogonal to each irreducible \mathbf{T} -module other than V_0 [6, Lem. 3.3]. For more information on V_0 see [6, 10]. We will use the following facts.

Lemma 4.1. [19, Lemma 3.9] *Let W denote an irreducible \mathbf{T} -module with endpoint r and diameter d . Then*

$$E_i^* W \neq 0 \quad (r \leq i \leq r + d). \quad (4.1)$$

Moreover

$$E_i^* A E_j^* W \neq 0 \quad \text{if } |i - j| = 1, \quad (r \leq i, j \leq r + d). \quad (4.2)$$

Lemma 4.2. [6, Lemma 3.4] *Let W denote a \mathbf{T} -module. Suppose there exists an integer i ($0 \leq i \leq D$) such that $\dim(E_i^*W) = 1$ and $W = \mathbf{T}E_i^*W$. Then W is irreducible.*

Theorem 4.3. [12, Lemma 10.1], [22, Theorem 11.1] *Let W denote a thin irreducible \mathbf{T} -module with endpoint one, and let v denote a nonzero vector in E_1^*W . Then $W = \mathbf{M}v$. Moreover the diameter of W is $D - 2$ or $D - 1$.*

Theorem 4.4. [12, Corollary 8.6, Theorem 9.8] *Let v denote a nonzero vector in E_1^*V which is orthogonal to s_1 . Then the dimension of $\mathbf{M}v$ is $D - 1$ or D . Suppose the dimension of $\mathbf{M}v$ is $D - 1$. Then $\mathbf{M}v$ is a thin irreducible \mathbf{T} -module with endpoint 1 and diameter $D - 2$.*

5 The proof of Theorem 1.1

We now give a proof of Theorem 1.1.

Proof. ((i) \implies (ii)) We show $\mathbf{M}v$ is a thin irreducible \mathbf{T} -module with endpoint 1. By Theorem 4.4 the dimension of $\mathbf{M}v$ is either $D - 1$ or D . First assume the dimension of $\mathbf{M}v$ is equal to $D - 1$. Then by Theorem 4.4, $\mathbf{M}v$ is a thin irreducible \mathbf{T} -module with endpoint 1. Next assume the dimension of $\mathbf{M}v$ is equal to D . The space $(\mathbf{M}; v)$ contains J and has dimension at least 2, so there exists $P \in (\mathbf{M}; v)$ such that J, P are linearly independent. From the construction $Pv \in E_D^*V$. Observe $Pv \neq 0$; otherwise the dimension of $\mathbf{M}v$ is not D . The elements A_0, A_1, \dots, A_D form a basis for \mathbf{M} . Therefore the elements $A_0 + A_1 + \dots + A_i$ ($0 \leq i \leq D$) form a basis for \mathbf{M} . Apparently there exist complex scalars ρ_i ($0 \leq i \leq D$) such that $P = \sum_{i=0}^D \rho_i (A_0 + A_1 + \dots + A_i)$. Recall $J = \sum_{h=0}^D A_h$. Subtracting a scalar multiple of J from P if necessary, we may assume $\rho_D = 0$. We consider Pv from two points of view. On one hand we have $Pv \in E_D^*V$. Therefore $E_D^*Pv = Pv$ and $E_i^*Pv = 0$ for $0 \leq i \leq D - 1$. On the other hand using (3.5),

$$Pv = \sum_{i=0}^{D-1} \rho_i (E_{i+1}^*A_i v - E_i^*A_{i+1} v).$$

Combining these two points of view we find $Pv = \rho_{D-1} E_D^* A_{D-1} v$, $E_0^* A v = 0$, and

$$\rho_{i-1} E_i^* A_{i-1} v = \rho_i E_i^* A_{i+1} v \quad (1 \leq i \leq D - 1). \quad (5.1)$$

We mentioned $Pv \neq 0$; therefore $\rho_{D-1} \neq 0$ and $E_D^* A_{D-1}v \neq 0$. Applying Corollary 3.5(i) we find $E_i^* A_{i-1}v \neq 0$ for $1 \leq i \leq D$. We claim $E_i^* A_{i+1}v$ and $E_i^* A_{i-1}v$ are linearly dependent for $1 \leq i \leq D-1$. Suppose there exists an integer i ($1 \leq i \leq D-1$) such that $E_i^* A_{i+1}v$ and $E_i^* A_{i-1}v$ are linearly independent. Then $E_i^* A_{i+1}v \neq 0$. Applying Corollary 3.5(ii) we find $E_j^* A_{j+1}v \neq 0$ for $i \leq j \leq D-1$. Using these facts and (5.1) we routinely find $\rho_j = 0$ for $i \leq j \leq D-1$. In particular $\rho_{D-1} = 0$ for a contradiction. We have now shown $E_i^* A_{i+1}v$ and $E_i^* A_{i-1}v$ are linearly dependent for $1 \leq i \leq D-1$. Observe $\mathbf{M}v$ is spanned by the vectors

$$(A_0 + A_1 + \cdots + A_i)v \quad (0 \leq i \leq D-1).$$

By (3.5) and our above comments we find $\mathbf{M}v$ is contained in the span of

$$E_{i+1}^* A_i v \quad (0 \leq i \leq D-1). \quad (5.2)$$

Since $\mathbf{M}v$ has dimension D we find $\mathbf{M}v$ is equal to the span of (5.2). Apparently $\mathbf{M}v$ is a \mathbf{T} -module. Moreover $\mathbf{M}v$ is irreducible by Lemma 4.2. Apparently $\mathbf{M}v$ is thin with endpoint 1.

((ii) \implies (i)) We show $(\mathbf{M}; v)$ has dimension at least 2. Since $J \in (\mathbf{M}; v)$ it suffices to exhibit an element $P \in (\mathbf{M}; v)$ such that J, P are linearly independent. Let W denote a thin irreducible \mathbf{T} -module which has endpoint 1 and contains v . By Theorem 4.3 we have $W = \mathbf{M}v$; also by Theorem 4.3 the diameter of W is $D-2$ or $D-1$. First suppose W has diameter $D-2$. Then W has dimension $D-1$. Consider the map $\sigma : \mathbf{M} \rightarrow V$ which sends each element P to Pv . The image of \mathbf{M} under σ is $\mathbf{M}v$ and the kernel of σ is contained in $(\mathbf{M}; v)$. The image has dimension $D-1$ and \mathbf{M} has dimension $D+1$ so the kernel has dimension 2. It follows $(\mathbf{M}; v)$ has dimension at least 2. Next assume W has diameter $D-1$. In this case $E_D^* W \neq 0$ by (4.1). Since $W = \mathbf{M}v$ there exists $P \in \mathbf{M}$ such that Pv is a nonzero element in $E_D^* W$. Now $P \in (\mathbf{M}; v)$. Observe P, J are linearly independent since $Pv \neq 0$ and $Jv = 0$. Apparently the dimension of $(\mathbf{M}; v)$ is at least 2.

Now assume (i), (ii) hold. We show the dimension of $(\mathbf{M}; v)$ is 2. To do this, we show the dimension of $(\mathbf{M}; v)$ is at most 2. Let H denote the subspace of \mathbf{M} spanned by A_0, A_1, \dots, A_{D-2} . We show H has 0 intersection with $(\mathbf{M}; v)$. By Theorem 4.4 the dimension of $\mathbf{M}v$ is at least $D-1$. Recall \mathbf{M} is generated by A so the vectors $A^i v$ ($0 \leq i \leq D-2$) are linearly independent. Apparently the vectors $A_i v$ ($0 \leq i \leq D-2$) are linearly independent. For $0 \leq i \leq D-2$

the vector $A_i v$ is contained in $\sum_{h=0}^{D-1} E_h^* V$ by Lemma 3.1(ii); therefore $A_i v$ is orthogonal to $E_D^* V$. We now see the vectors $A_i v$ ($0 \leq i \leq D-2$) are linearly independent and orthogonal to $E_D^* V$. It follows H has 0 intersection with $(\mathbf{M}; v)$. Observe H is codimension 2 in \mathbf{M} so the dimension of $(\mathbf{M}; v)$ is at most 2. We conclude the dimension of $(\mathbf{M}; v)$ is 2. \square

6 Pseudo primitive idempotents

In this section we introduce the notion of a pseudo primitive idempotent.

Definition 6.1. For each $\theta \in \mathbb{C} \cup \infty$ we define a subspace of \mathbf{M} which we call $\mathbf{M}(\theta)$. For $\theta \in \mathbb{C}$, $\mathbf{M}(\theta)$ consists of those elements Y of \mathbf{M} such that $(A - \theta I)Y \in \mathbb{C}A_D$. We define $\mathbf{M}(\infty) = \mathbb{C}A_D$.

With reference to Definition 6.1, we will show each $\mathbf{M}(\theta)$ has dimension 1. To establish this we display a basis for $\mathbf{M}(\theta)$. We will use the following result.

Lemma 6.2. Let Y denote an element of \mathbf{M} and write $Y = \sum_{i=0}^D \rho_i A_i$. Let θ denote a complex number. Then the following (i), (ii) are equivalent.

$$(i) \quad (A - \theta I)Y \in \mathbb{C}A_D.$$

$$(ii) \quad \rho_i = \rho_0 f_i(\theta) k_i^{-1} \text{ for } 0 \leq i \leq D.$$

Proof. Evaluating $(A - \theta I)Y$ using $Y = \sum_{i=0}^D \rho_i A_i$ and simplifying the result using (2.5) we obtain

$$(A - \theta I)Y = \sum_{i=0}^D A_i (c_i \rho_{i-1} + a_i \rho_i + b_i \rho_{i+1} - \theta \rho_i),$$

where $\rho_{-1} = 0$ and $\rho_{D+1} = 0$. Observe by (2.4), (2.6) that $\rho_i = \rho_0 f_i(\theta) k_i^{-1}$ for $0 \leq i \leq D$ if and only if $c_i \rho_{i-1} + a_i \rho_i + b_i \rho_{i+1} = \theta \rho_i$ for $0 \leq i \leq D-1$. The result follows. \square

Corollary 6.3. For $\theta \in \mathbb{C}$ the following is a basis for $\mathbf{M}(\theta)$.

$$\sum_{i=0}^D f_i(\theta) k_i^{-1} A_i. \tag{6.1}$$

Proof. Immediate from Lemma 6.2. \square

Corollary 6.4. *The space $\mathbf{M}(\theta)$ has dimension 1 for all $\theta \in \mathbb{C} \cup \infty$.*

Proof. Suppose $\theta = \infty$. Then $\mathbf{M}(\theta)$ has basis A_D and therefore has dimension 1. Suppose $\theta \in \mathbb{C}$. Then $\mathbf{M}(\theta)$ has dimension 1 by Corollary 6.3. \square

Lemma 6.5. *Let θ and θ' denote distinct elements of $\mathbb{C} \cup \infty$. Then $\mathbf{M}(\theta) \cap \mathbf{M}(\theta') = 0$.*

Proof. This is a routine consequence of Corollary 6.3 and the fact that $\mathbf{M}(\infty) = \mathbb{C}A_D$. \square

Corollary 6.6. *For $0 \leq i \leq D$ we have $\mathbf{M}(\theta_i) = \mathbb{C}E_i$.*

Proof. Observe $(A - \theta_i I)E_i = 0$ so $E_i \in \mathbf{M}(\theta_i)$. The space $\mathbf{M}(\theta_i)$ has dimension 1 by Corollary 6.4 and E_i is nonzero so E_i is a basis for $\mathbf{M}(\theta_i)$. \square

Remark 6.7. [2, p. 63] *For $0 \leq j \leq D$ we have*

$$E_j = m_j |X|^{-1} \sum_{i=0}^D f_i(\theta_j) k_i^{-1} A_i,$$

where m_j denotes the rank of E_j .

Definition 6.8. Let $\theta \in \mathbb{C} \cup \infty$. By a *pseudo primitive idempotent* for θ we mean a nonzero element of $\mathbf{M}(\theta)$, where $\mathbf{M}(\theta)$ is from Definition 6.1.

7 The local eigenvalues

Definition 7.1. Define a function $\tilde{\cdot} : \mathbb{C} \cup \infty \longrightarrow \mathbb{C} \cup \infty$ by

$$\tilde{\eta} = \begin{cases} \infty & \text{if } \eta = -1, \\ -1 & \text{if } \eta = \infty, \\ -1 - \frac{b_1}{1+\eta} & \text{if } \eta \neq -1, \eta \neq \infty. \end{cases}$$

Observe $\tilde{\tilde{\eta}} = \eta$ for all $\eta \in \mathbb{C} \cup \infty$.

Let v denote a nonzero vector in E_1^*V which is orthogonal to s_1 . Assume v is an eigenvector for $E_1^*AE_1^*$ and let η denote the corresponding eigenvalue. We recall a few facts concerning η and $\tilde{\eta}$. We have $\tilde{\theta}_1 \leq \eta \leq \tilde{\theta}_D$ [18, Theorem 1]. If $\eta = \tilde{\theta}_1$ then $\tilde{\eta} = \theta_1$. If $\eta = \tilde{\theta}_D$ then $\tilde{\eta} = \theta_D$. We have $\theta_D < -1 < \theta_1$ by [18, Lemma 3] so $\tilde{\theta}_1 < -1 < \tilde{\theta}_D$. If $\tilde{\theta}_1 < \eta < -1$ then $\theta_1 < \tilde{\eta}$. If $-1 < \eta < \tilde{\theta}_D$ then $\tilde{\eta} < \theta_D$. We will show that if $\tilde{\theta}_1 < \eta < \tilde{\theta}_D$ then $\tilde{\eta}$ is not an eigenvalue of Γ . Given the above inequalities, to prove this it suffices to prove the following result.

Proposition 7.2. *Let v denote a nonzero vector in E_1^*V . Assume v is an eigenvector for $E_1^*AE_1^*$ and let η denote the corresponding eigenvalue. Then $\tilde{\eta} \neq k$.*

Proof. Suppose $\tilde{\eta} = k$. Then $\eta = \tilde{k}$ so by Definition 7.1,

$$\eta = -1 - \frac{b_1}{k+1}.$$

By this and since $b_1 < k$ we see η is a rational number such that $-2 < \eta < -1$. In particular η is not an integer. Observe η is an eigenvalue of the subgraph of Γ induced on the set of vertices adjacent x ; therefore η is an algebraic integer. A rational algebraic integer is an integer so we have a contradiction. We conclude $\tilde{\eta} \neq k$. \square

Corollary 7.3. *Let v denote a nonzero vector in E_1^*V which is orthogonal to s_1 . Assume v is an eigenvector for $E_1^*AE_1^*$ and let η denote the corresponding eigenvalue. Suppose $\tilde{\theta}_1 < \eta < \tilde{\theta}_D$. Then $\tilde{\eta}$ is not an eigenvalue of Γ .*

8 The proof of Theorem 1.2

We now give a proof of Theorem 1.2.

Proof. We first show E is contained in $(\mathbf{M}; v)$. To do this we show $Ev \in E_D^*V$. First suppose $\eta \neq -1$. Then $\tilde{\eta} \in \mathbb{C}$ by Definition 7.1. By Definition 6.1 there exists $\epsilon \in \mathbb{C}$ such that $(A - \tilde{\eta}I)E = \epsilon A_D$. By this and Lemma 3.1(ii),

$$\begin{aligned} AEv &= \tilde{\eta}Ev + \epsilon A_D v \\ &\in \mathbb{C}Ev + E_{D-1}^*W + E_D^*W. \end{aligned} \tag{8.1}$$

In order to show $Ev \in E_D^*V$ we show $E_i^*Ev = 0$ for $0 \leq i \leq D-1$. Observe $E_0^*Ev = 0$ since $E_0^*Ev \in E_0^*W$ and W has endpoint 1. We show $E_1^*Ev = 0$. By Corollary 6.3 there exists a nonzero $m \in \mathbb{C}$ such that

$$E = m \sum_{h=0}^D f_h(\tilde{\eta}) k_h^{-1} A_h.$$

Let us abbreviate

$$\rho_h = m f_h(\tilde{\eta}) k_h^{-1} \quad (0 \leq h \leq D), \quad (8.2)$$

so that $E = \sum_{h=0}^D \rho_h A_h$. By this and (3.2) we find $E_1^*E E_1^* = \sum_{h=0}^2 \rho_h E_1^* A_h E_1^*$. Applying this to v we find

$$E_1^*Ev = \sum_{h=0}^2 \rho_h E_1^* A_h v. \quad (8.3)$$

Setting $i = 1$ in Lemma 3.1(i), applying each term to v , and using $Jv = 0$ we find

$$0 = \sum_{h=0}^2 E_1^* A_h v. \quad (8.4)$$

By (8.3), (8.4), and since $E_1^*Av = \eta v$ we find $E_1^*Ev = \gamma v$ where $\gamma = \rho_0 - \rho_2 + \eta(\rho_1 - \rho_2)$. Evaluating γ using (2.6), (8.2), and Definition 7.1 we routinely find $\gamma = 0$. Apparently $E_1^*Ev = 0$. We now show $E_i^*Ev = 0$ for $2 \leq i \leq D-1$. Suppose there exists an integer j ($2 \leq j \leq D-1$) such that $E_j^*Ev \neq 0$. We choose j minimal so that

$$E_i^*Ev = 0 \quad (0 \leq i \leq j-1). \quad (8.5)$$

Combining this with (8.1) we find

$$E_i^*AEv = 0 \quad (0 \leq i \leq j-1). \quad (8.6)$$

Since W is thin and since $E_j^*Ev \neq 0$ we find E_j^*Ev is a basis for E_j^*W . Apparently $E_{j-1}^*AE_j^*Ev$ spans $E_{j-1}^*AE_j^*W$. The space $E_{j-1}^*AE_j^*W$ is nonzero by (4.2) and since the diameter of W is at least $D-2$. Therefore $E_{j-1}^*AE_j^*Ev \neq 0$. We may now argue

$$\begin{aligned} E_{j-1}^*AEv &= \sum_{i=0}^D E_{j-1}^*AE_i^*Ev \\ &= E_{j-1}^*AE_j^*Ev && \text{by (3.3), (8.5)} \\ &\neq 0 \end{aligned}$$

which contradicts (8.6). We conclude $E_i^*Ev = 0$ for $2 \leq i \leq D-1$. We have now shown $E_i^*Ev = 0$ for $0 \leq i \leq D-1$ so $Ev \in E_D^*V$ in the case $\eta \neq -1$. Next suppose $\eta = -1$, so that $\tilde{\eta} = \infty$. By Definition 6.1 there exists a nonzero $t \in \mathbb{C}$ such that $E = tA_D$. In order to show $Ev \in E_D^*V$ we show $A_Dv \in E_D^*V$. Since A_Dv is contained in $E_{D-1}^*V + E_D^*V$ by Lemma 3.1(ii), it suffices to show $E_{D-1}^*A_Dv = 0$. To do this it is convenient to prove a bit more, that $E_i^*A_{i+1}^*v = 0$ for $1 \leq i \leq D-1$. We prove this by induction on i . First assume $i = 1$. Setting $i = 1$ in Lemma 3.1(i), applying each term to v and using $Jv = 0$, $E_1^*Av = -v$, we obtain $E_1^*A_2^*v = 0$. Next suppose $i \geq 2$ and assume by induction that $E_{i-1}^*A_iv = 0$. We show $E_i^*A_{i+1}^*v = 0$. To do this we assume $E_i^*A_{i+1}^*v \neq 0$ and get a contradiction. Note that $E_i^*A_{i+1}^*v$ spans E_i^*W since W is thin. Then $E_{i-1}^*AE_i^*A_{i+1}^*v \neq 0$ by (4.2). But $E_{i-1}^*AE_i^*A_{i+1}^*v = b_iE_{i-1}^*A_iv$ by Lemma 3.4(ii). Of course $b_i \neq 0$ so $E_{i-1}^*A_iv \neq 0$, a contradiction. Therefore $E_i^*A_{i+1}^*v = 0$. We have now shown $E_i^*A_{i+1}^*v = 0$ for $1 \leq i \leq D-1$ and in particular $E_{D-1}^*A_Dv = 0$. It follows $Ev \in E_D^*V$ for the case $\eta = -1$. We have now shown $Ev \in E_D^*V$ for all cases so $E \in (\mathbf{M}; v)$. We now prove E, J form a basis for $(\mathbf{M}; v)$. By Theorem 1.1 $(\mathbf{M}; v)$ has dimension 2. We mentioned earlier $J \in (\mathbf{M}; v)$. We show E, J are linearly independent. Recall E, J are pseudo primitive idempotents for $\tilde{\eta}, k$ respectively. We have $\tilde{\eta} \neq k$ by Proposition 7.2 so E, J are linearly independent in view of Lemma 6.5. \square

Acknowledgements The initial work for this paper was done when the second author was an Honorary Fellow at the University of Wisconsin-Madison (July-December 2000) supported by the National Science Council, Taiwan ROC.

References

- [1] J. M. Balmaceda and M. Oura. The Terwilliger algebras of the group association schemes of S_5 and A_5 . *Kyushu J. Math.*, 48(2):221–231, 1994.
- [2] E. Bannai and T. Ito. *Algebraic Combinatorics I: Association Schemes*. Benjamin/Cummings, London, 1984.
- [3] E. Bannai and A. Munemasa. The Terwilliger algebras of group association schemes. *Kyushu J. Math.*, 49(1):93–102, 1995.
- [4] A. E. Brouwer, A. M. Cohen, and A. Neumaier. *Distance-Regular Graphs*. Springer-Verlag, Berlin, 1989.

- [5] J. Caughman. The Terwilliger algebras of bipartite P - and Q - polynomial association schemes. *Discrete Math.*, 196:65–95, 1999.
- [6] B. Curtin. Bipartite distance-regular graphs I. *Graphs Combin.*, 15(2):143–158, 1999.
- [7] B. Curtin. Bipartite distance-regular graphs II. *Graphs Combin.*, 15(4):377–391, 1999.
- [8] B. Curtin. Distance-regular graphs which support a spin model are thin. *16th British Combinatorial Conference (London, 1997)*. *Discrete Math.* 197/198:205–216, 1999.
- [9] B. Curtin and K. Nomura. Spin models and strongly hyper-self-dual Bose-Mesner algebras. *J. Algebraic Combin.*, 13(2):173–186, 2001.
- [10] E. Egge. A generalization of the Terwilliger algebra. *J. Algebra*, 233:213–252, 2000.
- [11] E. Egge. The generalized Terwilliger algebra and its finite dimensional modules when $d = 2$. *J. Algebra*, 250:178–216, 2002.
- [12] J. T. Go and P. Terwilliger. Tight distance-regular graphs and the subconstituent algebra. *European J. Combin.*, to appear.
- [13] S. A. Hobart and T. Ito. The structure of nonthin irreducible T -modules: ladder bases and classical parameters. *J. Alg. Combin.*, 7:53–75, 1998.
- [14] H. Ishibashi, T. Ito, and M. Yamada. Terwilliger algebras of cyclotomic schemes and Jacobi sums. *European J. Combin.*, 20(5):397–410, 1999.
- [15] H. Ishibashi. The Terwilliger algebras of certain association schemes over the Galois rings of characteristic 4. *Graphs Combin.*, 12(1):39–54, 1996.
- [16] B. Sagan and J. S. Caughman. The multiplicities of a dual-thin Q -polynomial association scheme, *Electron. J. Combin.*, 8(1):(electronic) 2001.
- [17] K. Tanabe. The irreducible modules of the Terwilliger algebras of Doob schemes. *J. Alg. Combin.*, 6:173–195, 1997.
- [18] P. Terwilliger. A new feasibility condition for distance-regular graphs. *Discrete Math.*, 61:311–315, 1986.
- [19] P. Terwilliger. The subconstituent algebra of an association scheme I. *Journal of Algebraic Combinatorics*, 1(4):363–388, 1992.

- [20] P. Terwilliger. The subconstituent algebra of an association scheme II. *Journal of Algebraic Combinatorics*, 2(1):73-103, 1993.
- [21] P. Terwilliger. The subconstituent algebra of an association scheme III. *Journal of Algebraic Combinatorics*, 2(2):177-210, 1993.
- [22] P. Terwilliger. The subconstituent algebra of a distance-regular graph; thin modules with endpoint one. *Linear alg. Appl.*, to appear.
- [23] P. Terwilliger. An inequality involving the local eigenvalues of a distance-regular graph. *J. Algebraic Combin.*, submitted.
- [24] M. Tomiyama and N. Yamazaki. The subconstituent algebra of a strongly regular graph. *Kyushu J. Math.*, 48(2):323-334, 1994.

Paul Terwilliger
 Department of Mathematics
 University of Wisconsin
 480 Lincoln Drive
 Madison Wisconsin
 USA 53706
 Email: terwilli@math.wisc.edu

Chih-wen Weng
 Department of Applied Mathematics
 National Chiao Tung University
 1001 Ta Hsueh Road
 Hsinchu
 Taiwan ROC
 Email: weng@math.nctu.edu.tw